

May, 2001
OU-HET 387,
YITP-01-46,
hep-th/0105247

Some Computation on g Function and Disc Partition Function and Boundary String Field Theory

A. Fujii^{1*} and H. Itoyama^{2†}

¹*Yukawa Institute for Theoretical Physics, Kyoto University,
Kyoto 606-8502, Japan*

²*Department of Physics, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560-0043, Japan*

Abstract

Quadratic tachyon profile has been discussed in the boundary string field theory. We here compute the g -function by factorizing the cylinder amplitude. The answer is compared with the disc partition function. The boundary state is constructed. We extend these computations to those of the boundary sine-Gordon model at the free fermion point.

*e-mail address: fujii@yukawa.kyoto-u.ac.jp

†e-mail address: itoyama@het.phys.sci.osaka-u.ac.jp

The boundary string field (BSFT) theory[1] has been recently adopted for the analysis of Sen’s tachyon condensation[2] both in bosonic and supersymmetric string theory. In the bosonic BSFT the spacetime string action S is conjectured to satisfy [3]-[7]

$$\frac{\partial S}{\partial \lambda_i} = \mathcal{G}^{ij} \hat{\beta}_j, \quad (1)$$

where λ_i are the boundary coupling constants, $\mathcal{G}^{ij}$ is their metric, and $\hat{\beta}_i$ are the beta functions. This differential equation for S has the same form as that conjectured for the ground state degeneracy (g -function)[8] in statistical models. In fact the studies based on this similarity have been progressed by several authors [4][9][10].

In this letter we consider the model in which the open string on-shell action is deformed at the boundary by the quadratic term in X [1]. This model has been used in the evaluation of the exact tachyon potential [5]-[7] and can be obtained as a limit of boundary sine-Gordon (BSG) model. The Lagrangian of the BSG model is

$$\begin{aligned} \mathcal{S} &= \frac{1}{4\pi\alpha'} \int_{\Sigma} \sqrt{\det \eta} \eta^{ij} \partial_i X \partial_j X d^2\sigma + \int_{\partial\Sigma} T(X) dy, \\ T(X) &= \zeta \cos\left(\frac{2\pi}{R} X\right), \end{aligned} \quad (2)$$

where Σ is a worldsheet of open string, η_{ij} is its metric, and $\partial\Sigma$ is the boundary parameterized by y . If the quantity $\lambda = R^2/4\pi^2\alpha' - 1$ is a non-negative integer, the model is exactly solvable [11] and λ corresponds to the number of the species of the breathers. If we take the limit $R \rightarrow \infty$ and let the parameter ζ scale appropriately, $T(X)$ becomes a quadratic function of X , namely, $T(X) = a + uX^2$. Let us call this the boundary quadratic deformation (BQD) model.

We evaluate the partition functions of the BQD model on a cylinder and a disc. The disc partition function followed by the spacetime string action is already evaluated in ref.[1]. Our main interest in this letter goes to the g -description. Thus we first calculate the cylinder partition function and extract g by means of the finite size scaling. Because the BQD is a specific case of the BSG model with $R \rightarrow \infty$ limit, the direct calculation of the functional determinant is more suitable than the thermodynamic Bethe ansatz (TBA) method[12] [13], which includes breathers of an infinite number of species.

Secondly we construct the boundary state[14] on a disk which reproduces the known partition function[1] and can be used to define the g -function on the disk. We will see that S as a function of the boundary coupling constant on the disc has a pole at every minus integer whereas g on the cylinder has a zero there.

Finally we consider the BSG model at the free fermion (FF) point, $4\pi^2\alpha'/R^2 = 1/2$. The g -function of the FF model on a cylinder is obtained in refs.[13][15][17]. We evaluate its

partition function on a disc by means of the fermionization method. We find a similar result to the BQD case: g on the cylinder has a zero at every minus half-integer while the partition function on the disc possesses a pole there.

Let us consider the BQD model on a cylinder and evaluate its partition function. We remark that a similar analysis on an annulus can be found in ref.[19]. Let the length of the cylinder be l and the radius be r . The temperature is identified with $\theta = 1/2\pi r$. The Lagrangian and the partition function of the BQD model are

$$L = \frac{1}{4\pi\alpha'} \left[\int_0^l dx (\partial_a X)^2 - v X(t, 0)^2 \right] \quad (3)$$

and

$$Z = \int \mathcal{D}X \exp \left(- \int_0^{2\pi r} L dt \right), \quad (4)$$

respectively. The boundary condition at $x = 0$ is a mixed type:

$$\partial_x X + v X = 0. \quad (5)$$

We impose the Dirichlet type condition at $x = l$ and the periodic boundary condition in t -direction.¹

Let us calculate the partition function in the Lagrange formalism. We expand the field X as

$$X(t, x) = \sum_{m \in \mathbb{Z}} \sum_j \mathcal{A}_{m, \mu_j} e^{imt/r} \sin \mu_j (x - l), \quad \mathcal{A}_{m, \mu_j} = \mathcal{A}_{-m, \mu_j}. \quad (6)$$

Because we assume the periodic boundary condition in t -direction, m runs over integers. In x -direction, we impose the mixed boundary condition (5), which gives the equation determining μ_j :

$$\mu_j = v \tan(\mu_j l). \quad (7)$$

Therefore, the partition function becomes

$$Z' = \text{Det}'^{-1/2}(-\partial_t^2 - \partial_x^2) = \left(\left(\prod_{m=0}^{\infty} \prod_j \right)' [(m/r)^2 + \mu_j^2] \right)^{-1/2}. \quad (8)$$

The complete partition function Z includes the contribution from the zero mode integration.

Applying the standard zeta-function regularization, we obtain

$$Z' = \prod_{\mu_j > 0} (e^{\pi r \mu_j} - e^{-\pi r \mu_j})^{-1}. \quad (9)$$

¹This boundary condition is similar to but distinct from that of $p - p'$ open string with B -field [20][21].

Let us define a function

$$f(p) = p - \frac{1}{l} \tan^{-1} \frac{p}{v}. \quad (10)$$

If a certain μ_j satisfies (7), one can see $f(\mu_j) = \pi I_j / l$, where I_j is an integer. If $vl > 1$, $f(p)$ is a monotonically increasing function. Therefore, $I_{j+1} - I_j = 1$. Let us introduce the density function

$$\rho(\mu_j) = \lim_{l \rightarrow \infty} \frac{1}{l(\mu_{j+1} - \mu_j)}. \quad (11)$$

In the limit $l \rightarrow \infty$, μ_j develops continuously in $(0, \infty)$. From the form of (10), we see

$$\pi \rho(\mu) = 1 - \frac{1}{vl} \frac{1}{1 + (\mu/v)^2} + O(l^{-2}). \quad (12)$$

Hence, the free energy is represented like

$$\begin{aligned} F'/\theta = -\ln Z' &= \sum_{\mu_j > 0} [\pi r \mu_j + \ln(1 - e^{-2\pi r \mu_j})], \\ &= \int_0^\infty [\pi r \mu + \ln(1 - e^{-2\pi r \mu})] l \rho(\mu) d\mu. \end{aligned} \quad (13)$$

Let us recall the definition of g -factor in terms of the finite size scaling:

$$F = f_0 l - \theta \ln g + O(l^{-1}). \quad (14)$$

We conclude

$$\ln g = \frac{v}{\pi} \int_0^\infty \frac{d\mu}{\mu^2 + v^2} \ln(1 - e^{-2\pi r \mu}) - \ln \sqrt{rv} = \ln \left[\frac{\sqrt{2\pi}}{\Gamma(u+1)} \left(\frac{u}{e}\right)^u \right], \quad (15)$$

where $u = rv$. We have subtracted $\ln \sqrt{rv}$, taking the zero mode integration into account. In (15), taking the limits $u \rightarrow 0$ (UV limit) and $u \rightarrow \infty$ (IR limit), we see $g_{\text{UV}}/g_{\text{IR}} = \infty$. This is consistent with the g -theorem[8]. We should remark here that from (15) g has zeros at $u = -1, -2, \dots$.

Let us consider BQD model on a disc $\mathcal{D}$: $0 \leq r \leq c$, $0 \leq \sigma \leq 2\pi$, where c is the radius of $\mathcal{D}$. The action is

$$\mathcal{S} = \frac{1}{4\pi\alpha'} \int_{\mathcal{D}} \sqrt{\det \eta} \eta^{ij} g_{\mu\nu} \partial_i X^\mu \partial_j X^\nu d^2\sigma + \frac{1}{4\pi\alpha'} \int_{\partial\mathcal{D}} (a + u_{\mu\nu} X^\mu X^\nu) d\sigma, \quad (16)$$

where we set $\sigma_1 = r$ and $\sigma_2 = \sigma$. η^{ij} and $g_{\mu\nu}$ are the metrics of the world sheet $\mathcal{D}$ and the spacetime, respectively, a and $u_{\mu\nu}$ are the coupling constants at the boundary. From the action (16), we should impose the boundary condition on $\partial\mathcal{D}$ as

$$g_{\mu\nu} r \frac{\partial}{\partial r} X^\nu + u_{\mu\nu} X^\nu \Big|_{r=c} = 0. \quad (17)$$

In ref.[1] the partition function of the BQD model is calculated on the disc with unit radius, *i.e.* $c = 1$ with $g_{\mu\nu} = \delta_{\mu\nu}$ and $u_{\mu\nu} = u_\mu \delta_{\mu\nu}$. The result is

$$Z = e^{-a} \prod_{\mu} Z_1(u_{\mu}), \quad Z_1(u) = \sqrt{u} e^{\gamma u} \Gamma(u). \quad (18)$$

Let us consider the partition function (18) in the closed string picture. For this sake, we introduce the boundary state satisfying

$$\langle \mathcal{B} | \left(g_{\mu\nu} r \frac{\partial}{\partial r} X^\nu + u_{\mu\nu} X^\nu \right) \Big|_{r=c} = 0. \quad (19)$$

To determine the explicit form of $\langle \mathcal{B} |$, we utilize the mode expansion of X in terms of the complex coordinates, $z = r e^{i\sigma}$ and $\bar{z} = r e^{-i\sigma}$:

$$\begin{aligned} X^\mu(r, \sigma) &= X_R^\mu(z) + X_L^\mu(\bar{z}), \\ X_R^\mu(z) &= \frac{1}{2} X_0^\mu - \frac{i\alpha'}{2} p^\mu \ln z + i \sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \alpha_n^\mu \frac{z^{-n}}{n}, \\ X_L^\mu(\bar{z}) &= \frac{1}{2} X_0^\mu - \frac{i\alpha'}{2} p^\mu \ln \bar{z} + i \sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \tilde{\alpha}_n^\mu \frac{\bar{z}^{-n}}{n} \end{aligned} \quad (20)$$

with

$$[X_0^\mu, p^\nu] = i g^{\mu\nu}, \quad [\alpha_m^\mu, \alpha_n^\nu] = m g^{\mu\nu} \delta_{m+n,0}, \quad [\tilde{\alpha}_m^\mu, \tilde{\alpha}_n^\nu] = m g^{\mu\nu} \delta_{m+n,0}, \quad [\alpha_m^\mu, \tilde{\alpha}_n^\nu] = 0. \quad (21)$$

With the above expression, let us seek for the boundary state of the following form

$$\langle \mathcal{B} | = \langle 0 | \exp \left(-\frac{1}{2} X_0^\mu A_{\mu\nu} X_0^\nu \right) \exp \left(\sum_{m=1}^{\infty} \tilde{\alpha}_m^\mu C_{\mu\nu}^{(m)} \alpha_m^\nu \right), \quad (22)$$

where $\langle 0 |$ is the out vacuum state:

$$\langle 0 | \alpha_{-n}^\mu = \langle 0 | \tilde{\alpha}_{-n}^\mu = 0 \quad \text{for } n > 0. \quad (23)$$

Upon setting $g_{\mu\nu} = \delta_{\mu\nu}$ and $u_{\mu\nu} = u_\mu \delta_{\mu\nu}$, we see that the boundary state (22) satisfies the condition (19) if the parameters are

$$A_{\mu\nu} = \frac{1}{\alpha'} \frac{u_\mu}{1 + u_\mu \ln c} \delta_{\mu\nu}, \quad C_{\mu\nu}^{(m)} = -\frac{c^{-2m}}{m} \frac{m - u_\mu}{m + u_\mu} \delta_{\mu\nu}. \quad (24)$$

This boundary state (22) with (24) can be used in the calculation of the correlation functions. For example, the two point function can be calculated as

$$\begin{aligned} \frac{2}{\alpha'} \langle X^\mu(z, \bar{z}) X^\nu(w, \bar{w}) \rangle &= \frac{2 \langle \mathcal{B} | X^\mu(z, \bar{z}) X^\nu(w, \bar{w}) | 0 \rangle}{\langle \mathcal{B} | 0 \rangle} \\ &= -\delta^{\mu\nu} \ln(|z - w|/c)^2 - \delta^{\mu\nu} \ln |1 - z\bar{w}/c^2|^2 \\ &\quad + \delta^{\mu\nu} \cdot \frac{2}{u_\mu} - 2\delta^{\mu\nu} u_\mu \sum_{m=1}^{\infty} \frac{1}{m(m + u_\mu)} [(z\bar{w}/c^2)^m + (\bar{z}w/c^2)^m], \end{aligned} \quad (25)$$

which agrees with the result in ref.[1] by means of Green's differential equation. Hence, the boundary state (22) with (24) reproduces the disc partition function (18) precisely. The partition function (18) and the spacetime action $S(=i_W Z)$ have poles at $u_\mu = -1, -2, \dots$.

Finally let us evaluate the partition function of the BSG on a disk at the FF point, namely,

$$\mathcal{S}_{\text{SG}} = \frac{1}{8\pi} \int_D d^2\sigma \partial X \bar{\partial} X + \zeta \int_{\partial D} d\sigma \cos \frac{1}{2} X, \quad (26)$$

where we set $\alpha' = 2$. To execute the calculation, it is useful to apply the description in terms of massless fermions [15]-[18] of the action (26). The fermionic action $\mathcal{S}$ consists of four terms, $\mathcal{S} = \mathcal{S}_{\text{free}}^{(0)} + \mathcal{S}_{\text{free}}^{(1)} + \mathcal{S}_{\text{int}}^{(0)} + \mathcal{S}_{\text{int}}^{(1)}$, where

$$\mathcal{S}_{\text{free}}^{(0)} = \frac{1}{8\pi} \int_0^c r dr \int_0^{2\pi} d\sigma [\psi_+ \bar{\partial} \psi_- + \psi_- \bar{\partial} \psi_+ - \bar{\psi}_+ \partial \bar{\psi}_- - \bar{\psi}_- \partial \bar{\psi}_+] \quad (27)$$

is a free fermion action in the bulk, and the boundary terms are

$$\mathcal{S}_{\text{free}}^{(1)} = -\frac{c}{8\pi} \int_0^{2\pi} d\sigma [\psi_+ \bar{\psi}_+ + \psi_- \bar{\psi}_-], \quad (28)$$

$$\mathcal{S}_{\text{int}}^{(0)} = \frac{1}{2\pi} \int_0^{2\pi} d\sigma a_+ \partial_\sigma a_-, \quad (29)$$

$$\mathcal{S}_{\text{int}}^{(1)} = \frac{c\zeta}{32\pi^2} \int_0^{2\pi} d\sigma [e^{i\sigma/2}(\psi_+ a_- + a_+ \psi_-) + e^{-i\sigma/2}(\bar{\psi}_- a_+ + a_- \bar{\psi}_+)]. \quad (30)$$

We can obtain the equation of motion

$$\bar{\partial} \psi_\pm = \partial \bar{\psi}_\pm = 0, \quad (31)$$

and the boundary conditions

$$a_+ = a_- = e^{i\sigma/2} \psi_+ - e^{-i\sigma/2} \bar{\psi}_- = -e^{i\sigma/2} \psi_- + e^{-i\sigma/2} \bar{\psi}_+ \quad (32)$$

$$0 = \partial_\sigma (e^{-i\sigma/2} \bar{\psi}_+ - e^{i\sigma/2} \psi_-) - B(e^{-i\sigma/2} \bar{\psi}_- - e^{i\sigma/2} \psi_+), \quad (33)$$

where dimensionless parameter is introduced, $B = \zeta^2 c / 64\pi^2$.

Substituting (32) into (30), we obtain

$$-\frac{d}{d\zeta} \ln Z = \frac{c}{2} \int_0^{2\pi} d\sigma [e^{i\sigma/2}(\langle \psi_+ a_- \rangle + \langle a_+ \psi_- \rangle) + e^{-i\sigma/2}(\langle \bar{\psi}_- a_+ \rangle + \langle a_- \bar{\psi}_+ \rangle)] \quad (34)$$

With the help of Green's functions, we can calculate the partition function by integrating (34). As a result, we should integrate

$$-\frac{d}{d\zeta} \ln Z = -\lim_{\epsilon \rightarrow 0} \frac{1}{\pi\zeta} \int_0^{2\pi} d\sigma \sum_{k=0}^{\infty} \frac{k+1/2}{k+1/2+B} e^{-ik\epsilon}. \quad (35)$$

The above quantity diverges as $\epsilon \rightarrow 0$, and thus needs regularization. If we subtract $1 - B/(k + 1/2)$ in the summation, we obtain

$$-\frac{d}{d\zeta} \ln Z_{\text{reg}} = -\frac{B^2}{\pi\zeta} \int_0^{2\pi} d\sigma \sum_{k=0}^{\infty} \frac{1}{(k + 1/2)(k + 1/2 + B)}. \quad (36)$$

We can integrate (36) to yield

$$Z_{\text{reg}} = \frac{\Gamma(2B)}{\Gamma(B)} = \frac{2^{2B-1}}{\sqrt{\pi}} \Gamma(B + 1/2), \quad (37)$$

which has poles at $B = -1/2, -3/2, \dots$. On the other hand, the g -function defined on a cylinder is known[13][15][17] as

$$g = \frac{\sqrt{2\pi}}{\Gamma(\alpha + 1/2)} \left(\frac{\alpha}{e}\right)^\alpha, \quad (38)$$

where $\alpha = 32\pi^2 r \zeta^2$.

We finally point out again a curious reciprocal relation between the disc partition function and the g -function extracted from the cylinder partition function, which is common to the two examples considered in this letter. In the analytic coupling constant plane, the disc partition function has integer spaced poles in the negative real axis whereas the g -function possesses zeros precisely at these points.

Acknowledgments

The authors would like to thank T. Nakatsu and T. Suyama for helpful discussions. This work is supported in part by the Grant-in-Aid for Scientific Research No.12640272 from the Ministry of Education, Science and Culture, Japan. AF was partially supported by The Yukawa Memorial Foundation.

References

- [1] E. Witten, Phys. Rev. **D46**(1992)5467, [hep-th/9208027](#); Phys. Rev. **D47**(1993)3405, [hep-th/9210065](#).
- [2] A. Sen, Int. J. Mod. Phys. **14**(1999)4061, [hep-th/9902105](#).
- [3] S. Shatashvili, Phys. Lett. **B311**(1993)83, [hep-th/9303143](#); “*On the problems with background independence in string theory*”, [hep-th/9311177](#).
- [4] J.A. Harvey, D. Kutasov and E. Martinec, “*On the relevance of tachyons*”, [hep-th/0003101](#).

- [5] A.A. Gerasimov and S.L. Shatashvili, JHEP **0010**(2000)034, [hep-th/0009103](#).
- [6] D. Kutasov, M. Mariño and G. Moore, JHEP **0010**(2000)045, [hep-th/0009148](#).
- [7] A.A. Tseytlin, “*Sigma model approach to string field theory effective actions with tachyons*”, To appear in special issue of J.Math.Phys. on Strings, Branes and M Theory, [hep-th/0011033](#).
- [8] I. Affleck and A.W. Ludwig, Phys. Rev. Lett. **67**(1991)161; Phys. Rev. **B48**(1993)7297.
- [9] J.A. Harvey, S. Kachru, G. Moore and E. Silverstein, JHEP **0003**(2000)001, [hep-th/9909072](#).
- [10] A. Fujii and H. Itoyama, Phys. Rev. Lett. **86**(2001)5235, [hep-th/0012150](#).
- [11] I.V. Cherednik, Theo. Math. Phys. **61**(1984)977; S. Ghoshal and A.B. Zamolodchikov, Int. J. Mod. Phys. **A9**(1994)3841, [hep-th/9306002](#).
- [12] P. Fendley, H. Saleur and N.P. Warner, Nucl. Phys. **B430**(1994)577, [hep-th/9406125](#).
- [13] A. LeClair, G. Mussardo, H. Saleur and S. Skorik, Nucl. Phys. **B453**(1995)581, [hep-th/9503227](#).
- [14] C.G. Callan, C. Lovelace, C.R. Nappi and S.A. Yost, Nucl. Phys. **B288** (1987) 525; A. Abouelsaood, C.G. Callan, C.R. Nappi and S.A. Yost, Nucl. Phys. **B280** (1987) 599; E.S. Fradkin and A.A. Tseytlin, Phys. Lett. **163B** (1985) 123.
- [15] R. Chatterjee, Mod. Phys. Lett. **A10**(1995)973, [hep-th/9412169](#).
- [16] M. Ameduri, R. Konik and A. LeClair, Phys. Lett. **B354**(1995)376, [hep-th/9503088](#).
- [17] R.M. Konik, “*Massless boundary sine-Gordon at the free fermion point: correlation and partition functions with applications to quantum wires*”, [hep-th/9507053](#).
- [18] L. Mezincescu and R.I. Nepomechie, ”*Integrals of motion for the sine-Gordon model with boundary at the free Fermion point*” in *Unified Symmetry: In the Small and in the Large 2*, ed. by B. Kursunoglu, S. Mintz and A. Perlmutter, Plenum Publishing Co., 213 - 219 (1995).
- [19] T. Suyama, “*Tachyon condensation and spectrum of strings on D-branes*”, [hep-th/0102192](#); K. Bardakci and A. Konechny, “*Tachyon condensation in boundary string field theory at one loop*”, [hep-th/0105098](#); T. Lee, “*Tachyon condensation, boundary state and noncommutative solitons*”, [hep-th/0105115](#).

- [20] N. Seiberg and E. Witten, JHEP **9812**(1998)008, hep-th/9908142.
- [21] B. Chen, H. Itoyama, T. Matsuo and K. Murakami, Nucl. Phys. **B576**(2000)177, hep-th/9910263; Nucl. Phys. **B593**(2001)505, hep-th/0005283; “*Correspondence between noncommutative soliton and open string/D-brane system via Gaussian damping factor*”, hep-th/0010066.